

42646

NOV 10 2008

REVISED
DATE REVIEWER

OCT -- 2008

Original ENGLISH FORMULA



CORRY'S

NOT AFFECTED BY RAIN

SLUG & SNAIL DEATH (3.25)

CORRY'S

SLUG & SNAIL DEATH (3.25)

PULL TAB TO OPEN

- Easy Application Container
- Fast, Effective Control Of Slugs and Snails
- Protects Fruit, Vegetable, Flower And Ornamental Gardens
- Contains Denatonium Benzoate

NOTICE:

To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.


Manufactured by
Matson, LLC. ©10-08
P. O. Box 1820

North Bend, Washington 98045
EPA Reg. No. 8119-11
EPA Est. No. 8119-WA-1
www.protectyourgarden.com
www.metaworksbest.com

META® is a registered trademark of Lonza Ltd. and the right to use the META mark has been granted to Matson, LLC. by Lonza Inc.

NOV 19 2008 8119-11 OR

IDEAL FOR VEGETABLE GARDENS



Easy To Use Meal Formula

KEEP OUT OF REACH OF CHILDREN

CAUTION: THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN. KEEP PETS OUT OF TREATED AREA. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS & FIRST AID.

NET.WT. 3.5 LB (1.58 kg)

Active Ingredient:
Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxocyclo-octane).....3.25%
Other Ingredients.....96.75%



**Can Be Used Around Fruit and Vegetable Plants,
Lawns and Ornamental Plants too!**

Provides Up To Two Weeks of Plant Protection.

Slugs and snails are two of our most destructive garden pests. Although their outward appearance is quite different, they are essentially the same creature, except that the slug does not have the external shell of the snail.

Slugs and snails feed almost entirely at night, in the evening or during dark days. They prefer moist, warm conditions and rarely venture from their hiding places during cold weather or during exceedingly dry or hot spells.

Slugs and snails hatch from eggs, laid in clusters, under debris, stones, compost heaps or beneath the surface of the soil. Generally, they will hatch out after 20 to 30 days of favorable weather conditions after which time they will continue to grow slowly, eating our most tender and succulent vegetation.

While these small slugs and snails are growing and eating they migrate towards better feeding and sheltering conditions. It is because of this migrating habit that it is impossible to completely eliminate the slug and snail problem. However, with the use of **Corry's® Slug & Snail Death (3.25)** and a consistent baiting program, these pests can be reduced to where they no longer present a major threat to our gardens.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO APPLY: (See graphics for suggested application methods) This product is for use in Home and Garden areas. **Corry's® Slug & Snail Death (3.25)** is so easy to use. This product requires no mixing, spraying or special applicators. Just scatter lightly on the soil surface in infested areas as recommended in the graphics below. Apply heavier if infestation is severe. If slugs and snails are invading from adjacent properties, a pencil thin line the entire length of the infested area will form a barrier resulting in excellent control. This bait is extremely attractive to slugs and snails and actually lures them from their hiding places and kills them. Reapply bait every two weeks – more often during heavy infestations or in areas heavily watered. All likely places should be treated as these pests migrate. They favor damp places around vegetable plants, flower gardens, rockeries, hedges, compost heaps, dichondra lawns, ivy patches and other ground cover where they can obtain shelter by day. For best results, be sure area to be baited is damp. In very dry weather, water well before baiting. Use approximately one pound per 2000 square feet.



CORRECT



CORRECT



CORRECT



INCORRECT



INCORRECT

Corry's® Slug & Snail Death (3.25) can be used to protect the following fruit and vegetable plants. Apply only to the soil surface. Do not apply to the foliage or edible parts of the plant. Use **Corry's® Slug & Snail Death (3.25)** to protect asparagus, beans, cabbage, carrots, celery, cucumbers, lettuce, melons, onions, peas, peppers, potatoes, radishes, spinach, squash, tomatoes, turnips, apples, avocados, blackberries, cherries, citrus, grapes, peaches, plums and strawberries.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry, locked location out of direct sunlight, in original container, and not accessible to children or pets. Avoid contamination of feed and foodstuffs.

PESTICIDE DISPOSAL: If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after use. **KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.** This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area.

IMPORTANT: Keep away from children, dogs and pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed. Use and store under conditions where there will not be a hazard.

FIRST AID

IF SWALLOWED:

- * Call poison control center or doctor immediately for treatment advice.
- * Have person sip a glass of water if able to swallow.
- * Do not induce vomiting unless told to do so by the poison control center or doctor.
- * Do not give anything by mouth to an unconscious person.

IF INHALED:

- * Move person to fresh air.
- * If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth, if possible.
- * Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING:

- * Take off contaminated clothing.
- * Rinse skin immediately with plenty of water for 15-20 minutes.
- * Call a poison control center or doctor for treatment advice.

IF IN EYES:

- * Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- * Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- * Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call 1-800-308-3723.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.