

Dimension® 0.125% Plus Fertilizer Turf and Ornamental Herbicide

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

This bag covers 11,350 sq ft

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, this product may be applied only by commercial applicators at no more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

| | |
|-------------------------------------------------------------------------------------------------------------------------------------------|---------|
| Active ingredient dithiopyr, 3,5-pyridinedicarbonyl acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester | 0.125% |
| Other ingredients | 99.875% |
| Total | 100.0% |

This product contains 0.0825 pound of the active ingredient dithiopyr per 50 pound bag

Keep Out of Reach of Children

CAUTION

| | | |
|------------------------------------------|--------------------------------------------|-------------------------------------------|
| EPA EST. NO. 87245-CAN-1 (Lot begins Z)* | EPA Reg. No. 62719-483-84886 | EPA EST. NO. 49991-TX-2 (Lot begins TX-2) |
| *If lot begins "Z", PRODUCT OF CANADA | EPA EST. NO. 75527-PA-12 (Lot ends PA-012) | EPA EST. NO. 89135-AZ-1 (Lot begins 4) |
| EPA EST. NO. 41124-W-1 (Lot begins W) | EPA EST. NO. 69759-OR-2 (Lot begins MAS) | EPA EST. NO. 35512-FL-2 (Lot ends GF) |
| EPA EST. NO. 57131-W-1 (Lot begins EC) | EPA EST. NO. 41124-OH-1 (Lot begins O) | EPA EST. NO. 82757-FL-1 (Lot begins S) |
| EPA EST. NO. 82757-OH-1 (Lot begins MF) | EPA EST. NO. 82757-MA-2 (Lot begins H) | |

Net Weight 50 LBS. (22.68 Kg)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation • Harmful If Inhaled

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, the give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. **Hot Line Number:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.

Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning, if burned, stay out of smoke.

Use Directions for Turf

This product provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

Use Precautions for Turf

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may increase the chances of turf injury.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

| Cool-Season Grasses | Warm-Season Grasses | |
|----------------------------------|---------------------|-----------------------------|
| bentgrass, creeping [†] | Agrostis palustris | bahiagrass |
| bluegrass, Kentucky | Poa pratensis | bermudagrass |
| fescue, fine ^{††} | Festuca rubra | buffalograss ^{†††} |
| fescue, tall | Festuca arundinacea | carpetgrass |
| ryegrass, perennial | Lolium perenne | centipedegrass |
| | | kikuyograss |
| | | St. Augustinegrass |
| | | zoysiagrass |
| | | Paspalum notatum |
| | | Cynodon dactylon |
| | | Buchloe dactyloides |
| | | Axonopus affinis |
| | | Eremochloa ophiuroides |
| | | Pennisetum clandestinum |
| | | Stenotaphrum secundatum |
| | | Zoysia japonica |

[†] Use of this product on certain varieties of creeping bentgrass, such as 'ochansey', 'carmen', 'seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colonial bentgrass (Agrostis tenuis).

^{††} Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'banner', 'beauty', 'bigart', 'CF-2', 'enjoy', 'HF-93', 'highlight', 'ivalo', 'Jamestown', 'kicket', 'majenta', 'mary', 'pennlawn', 'tamara', 'tajana', 'waldorf', and 'waldina'. Not all varieties of fine fescue have been tested.

^{†††} Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

Precautions for Reseeding, Overseeding, or Sprigging

- Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses.
- When reseeding or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation and fertilization.
- For best overseeding results, use mechanical or power seeding equipment (sill seeders) designed to provide good contact between soil and seed.

Precautions for Control of Crabgrass

Early postemergence weed control with this product is limited to crabgrass. Postemergence application effectively controls crabgrass, but only if application is made prior to the 3-leaf stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aeration, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

Other Use Precautions

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.
- Do not graze livestock or feed forage cut from areas treated with this product.

Application Directions for Turf

Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Use of split (half-rate) applications spaced 5 - 10 weeks apart may provide improved weed control.

Maximum Use Rates (Applies to all Turfgrass Uses)

- Do not apply more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per application or more than 1200 lb (1.5 lb active ingredient) per acre (27.6 lb per 1000 sq ft) per year if using split or sequential applications.
- In the state of New York, this product may be applied only by commercial applicators at no more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

32-0-8

Intermediate Size (SGN 190)

GUARANTEED ANALYSIS

| | |
|-----------------------------------------|--------|
| Total Nitrogen (N) | 32.00% |
| 32.00% Urea Nitrogen* | |
| Soluble Potash (K ₂ O) | 8.00% |

DERIVED FROM: Polymer Coated Urea, Urea and Sulfate of Potash

*26.8% Slowly Available Urea Nitrogen from Polymer Coated Urea
WARNING: THIS PRODUCT CONTAINS CHEMICALS KNOWN TO THE STATE OF CALIFORNIA TO CAUSE CANCER OR BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM. Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, 'Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant'" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/gfbmp07.pdf>.
 Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at http://www.dep.state.fl.us/central/Home/Meetings/Training/FLGreen/BMP_Book_final.pdf.

Product No. 8121021 F1828

GENERAL APPLICATION GUIDELINES

| Rate | lbs. Product per 1,000 sq ft | lbs. Nitrogen per 1,000 sq ft | lbs. A.I. per acre |
|--------|------------------------------|-------------------------------|--------------------|
| Low | 6.3 | 2.0 | .34 |
| Medium | 7.5 | 2.4 | .42 |
| High | 8.8 | 2.8 | .48 |



SPREAD IT & FORGET IT
 for your turf all season long

For Best Results use Quality Cut Turf

This product contains Duration CR, a registered trademark of Agrium, Inc.



GUARANTEED BY:
 Agrium Advanced Technologies (U.S.) Inc.
 2915 Rocky Mountain Avenue
 Loveland, CO 80538

Control of Crabgrass

Preemergence and Early Postemergence Control

- This product provides "preemergence" control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass.
- This product also provides "early postemergence" control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergence crabgrass control is a 2 - 4 week longer period of time (window of application) during which crabgrass control can be achieved.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described in the following use programs.

Spring Applications

For applications made in the spring or early summer, use an appropriate application rate corresponding to one of the three control programs listed in Table 1. Specific rate recommendations are based on the user's location (Region), the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where turfgrass is cut relatively high (e.g., homeowner lawns). This program provides 3 - 5 months of preemergence crabgrass control. This program will also provide early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., homeowner lawns).

Use Program 2 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., residential lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively low (e.g., golf fairways).

Table 1 Recommended Use Rates

| Region | Application Rates | Program 1 | Program 2 | Program 3 |
|-----------------------------------------------------------------------------------------------------------------------------|-------------------|-------------|-------------|-------------|
| All states, except NY, and parts of states not included in transition, south, coastal south or west. | lb/acre | 152 | 200 | 304 |
| | lb/1000 sq ft | 3.5 | 4.6 | 7.0 |
| | lb ai/acre | 0.19 | 0.25 | 0.38 |
| Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, & RI | lb/acre | 200 | 304 | 400 |
| | lb/1000 sq ft | 4.6 | 7.0 | 9.2 |
| | lb ai/acre | 0.25 | 0.38 | 0.5 |
| South: AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX | lb/acre | 304 | 200 + 200 | 248 + 248 |
| | lb/1000 sq ft | 7.0 | 4.6 + 4.6† | 5.7 + 5.7† |
| | lb ai/acre | 0.38 | 0.25 + 0.25 | 0.31 + 0.31 |
| Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX | lb/acre | 200 + 200 | 248 + 248 | 304 + 304 |
| | lb/1000 sq ft | 4.6 + 4.6† | 5.7 + 5.7† | 7.0 + 7.0† |
| | lb ai/acre | 0.25 + 0.25 | 0.31 + 0.31 | 0.38 + 0.38 |
| West: AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons. | lb/acre | 152 + 200 | 200 + 304 | 200 + 200 |
| | lb/1000 sq ft | 3.5 + 4.6 | 4.6 + 7.0 | 4.6 + 4.6† |
| | lb ai/acre | 0.19 + 0.25 | 0.25 + 0.38 | 0.25 + 0.25 |

† Split applications may be made 5 - 10 weeks apart.

Use of Split or Sequential Applications: Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with the rates in Table 1 being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application, provided the maximum application rate per year is not exceeded. Refer to the "Maximum Use Rates" section above and separate rate restrictions for the state of New York.

Fall Applications

This product can also be applied in the late summer through early fall (late August through November) at use rates listed in Table 1 for "Program 3" to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for seasonal control of crabgrass.

Weeds Controlled or Suppressed

Spring Applications

When applied preemergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds at application rates given in Table 1 above:

| Grasses | | | |
|------------------------|----------------------------------|-------------------------------|--------------------------------|
| barley | <i>Hordeum</i> spp. | zoosegrass | <i>Elymus indica</i> |
| barnyardgrass | <i>Echinochloa crus-galli</i> | foxtail, green | <i>Setaria verti</i> |
| bluegrass, annual | <i>Poa annua</i> | foxtail, yellow | <i>Setaria faberi</i> |
| brome | <i>Bromus</i> spp. | kikuyugrass † | <i>Pennisetum clandestinum</i> |
| crabgrass, large | <i>Digitaria sanguinalis</i> | oats, wild | <i>Avena fatua</i> |
| crabgrass, smooth | <i>Digitaria ischaemum</i> | ryegrass (annual & perennial) | <i>Lolium</i> spp. |
| crabgrass, Southern | <i>Digitaria ciliaris</i> | sandbur | <i>Cenchrus</i> spp. |
| crowfootgrass † | <i>Dactyloctenium aegyptium</i> | smutgrass | <i>Sporobolus indicus</i> |
| dallisgrass (seedling) | <i>Paspalum dilatatum</i> | | |
| Broadleaf Weeds | | | |
| bittercress † | <i>Cardamine</i> spp. | pigweed, redroot | <i>Amaranthus retroflexus</i> |
| carpetweed | <i>Mollugo verticillata</i> | parsley-piert † | <i>Alchemilla arvensis</i> |
| chickweed † | <i>Stellaria</i> spp. | purslane, common | <i>Portulaca oleracea</i> |
| geranium, Carolina † | <i>Geranium carolinianum</i> | rocket, London | <i>Sisymbrium irio</i> |
| henbit | <i>Lamium</i> spp. | shepherdspurse | <i>Capsella bursa-pastoris</i> |
| knotweed, prostrate | <i>Polygonum aviculare</i> | speedwell, corn † | <i>Veronica arvensis</i> |
| lespedeza, common † | <i>Lespedeza striata</i> | spurge, garden | <i>Euphorbia hirta</i> |
| marehail | <i>Conyza canadensis</i> | spurge, prostrate | <i>Euphorbia humifusa</i> |
| medic, black | <i>Medicago lupulina</i> | spurge, spotted | <i>Euphorbia maculata</i> |
| mustard | <i>Brassica</i> spp. | woodsorrel, creeping | <i>Oxalis corniculata</i> |
| oxalis, buttercup | <i>Oxalis pes-caprae</i> | woodsorrel, yellow | <i>Oxalis stricta</i> |
| pineappleweed † | <i>Matricaria matricarioides</i> | | |

† Suppression only

Fall Applications

When applied preemergence (prior to germination) in late summer through early fall (late August through November) this product will control or suppress the following weeds in addition to providing preemergence control of crabgrass in the following spring:

| | | | |
|-------------------|-----------------------|--------------------|----------------------------------|
| bluegrass, annual | <i>Poa annua</i> | geranium, Carolina | <i>Geranium carolinianum</i> |
| bittercress | <i>Cardamine</i> spp. | parsley-piert | <i>Alchemilla arvensis</i> |
| chickweed | <i>Stellaria</i> spp. | pineappleweed | <i>Matricaria matricarioides</i> |
| henbit | <i>Lamium</i> spp. | shepherdspurse | <i>Capsella bursa-pastoris</i> |

Application Instructions and Equipment

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

Use Directions for Landscape Ornamentals

This product provides preemergence control or suppression of listed annual grass and broadleaf weeds in plantings of ornamental plants listed on this label. This product may be applied in ornamental gardens, parks, golf courses and residential areas where ornamental plants are grown for aesthetic purposes. Apply this product with a properly calibrated spreader that will provide uniform particle distribution. Follow instructions in the "Application Instructions and Equipment" section above.

Use Rates

This product may be applied as a single application or split application. Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. To make a split application, divide the rate in the following table into separate applications made 5-10 weeks apart. Sequential applications may be made in the late summer following a spring application or in the spring after a fall application, provided maximum application rates per year given below are not exceeded.

| Pounds of Product/Acre | Pounds of Product/1000 sq ft | Pounds of Product/100 sq ft † |
|------------------------|------------------------------|-------------------------------|
| 400 | 9.2 | 0.92 (14.7 oz) |

† Rate based on an area of 100 sq ft for treatment of small areas.

Maximum Use Rates

- Do not apply more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per application or more than 1200 lb (1.5 lb active ingredient) per acre (27.6 lb per 1000 sq ft) per year if using split or sequential applications.
- In the state of New York, this product may be applied only by commercial applicators at no more than 400 lb (0.5 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Application Timing and Recommendations

This product is effective as a preemergence herbicide, but will not control established weeds. Applications to mulched areas or bare ground must be made prior to weed seed germination. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Existing weedy vegetation may be controlled by hand weeding, cultivation or using postemergence herbicides prior to application.

Tips for Improved Weed Control

- The herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Avoid disturbance of treated areas. Loss of weed control may result if the treated soil surface is disturbed by soil mixing or tillage.

Use Precautions

- Apply this product only to established ornamentals.
- Do not apply this product directly to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress resulting from drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides, or damage due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove any herbicide granules adhering to foliage.
- Do not apply under conditions that would affect uniformity of application or distribution on the soil surface. Uneven product distribution will result in uneven weed control. Application under windy conditions can result in uneven distribution or cause herbicide granules to drift from the intended treatment area.
- Do not graze livestock or feed forage cut from areas treated with this product.

Weeds Controlled or Suppressed

Used as directed, this product will control crabgrass and control or suppress other listed grass and broadleaf weeds when applied prior to their germination. Refer to "Use Directions for Turf" section for a complete listing of weeds controlled or suppressed. This product will not control established broadleaf weeds or grasses, except for crabgrass in early stages of development. The area to be treated should be free of weeds prior to application.

Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to directions in this label. However, this product has not been tested on all ornamental species, all cultivars of species on this label, or under all possible growing conditions. For species or cultivars not listed, it is recommended that a few plants in a limited area be treated and observed for tolerance under local growing conditions prior to large-scale use.

| Common Name | Botanical Name | Tolerant Cultivars |
|--------------------|----------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| abelia, dwarf | <i>Abelia X grandiflora</i> | none |
| ajuga | <i>Ajuga reptans</i> | bronze |
| | <i>Ajuga genevensis</i> | bronze beauty |
| almond, flowering | <i>Prunus glandulosa</i> | |
| apple † | <i>Malus pumila</i> | |
| arborvitae | <i>Thuja occidentalis</i> | nigra, pyramidalis, smaragd, techry, woodwards |
| arborvitae, golden | <i>Thuja orientalis</i> | |
| aster, Chinese | <i>Callistephus chinensis</i> | dwarf queen |
| ash, green | <i>Fraxinus pennsylvanica</i> | |
| ash, mountain | <i>Sorbus aucuparia</i> | |
| ash, purple | <i>Fraxinus americana</i> | |
| azalea | <i>Rhododendron</i> spp. | brilliant, buccaneer, caror, chimes (belgian), elise lee, exbury, fashion, harjoizer beauty, hershey red, higasa, hinoomison, holland (hybrid), marion lee, northern lights, orange cup, orchid lights, snow, southern charm |
| azalea, flame | <i>Rhododendron calendulacum</i> | |
| azalea, kirishima | | |
| bamboo, heavenly | | |
| barberry | <i>Berberis thunbergii</i> | aurae, dwarf pigmy, green, kobold, pigmy red, rose glow |
| barberry, purple | | isropurpurea |
| basket flower | <i>Gaillardia grandiflora</i> | |
| bearberry (common) | <i>Arctostaphylos uva-ursi</i> | Massachusetts |
| bee balm | <i>Monarda didyma</i> | |
| begonia | <i>Begonia</i> spp. | |
| birch, river | <i>Betula nigra</i> | |
| blackeyed Susan | <i>Rudbeckia hirta</i> | goldstrum |

| Common Name | Botanical Name | Tolerant Cultivars |
|-------------------------|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|
| blanket flower | Galathea spp. | |
| blueberry ¹ | Vaccinium spp. | bluecrop, blue jay, jersey, north blue, northland |
| bottlebrush | Callistemon citrinus | |
| boxwood, Japanese | Japonica | |
| boxwood, weiler | Buxis sempervirens | |
| broom | Cytisus spp. | moonlight |
| | Genista pilosa | Vancouver gold |
| bush carpet | | |
| camellia | Camellia japonica | debutante, mathollana supreme, chansonette |
| | Camellia sasanqua | |
| candy tuft | Iberis spp. | snow white |
| carex, variegated | Carex | |
| cedar, red | Juniperus virginiana | |
| celosia | Celosia spp. | |
| centaurea | Centaurea montana | |
| cockscorn, plumosa | Celosia cristata | scarlet plumosa |
| coleus | Coleus blumei | red kempia |
| columbine | Aquilegia spp. | |
| copper leaf | Acalypha wilkesiana | |
| coreopsis | Coreopsis spp. | moonbeam |
| corn flower | Centauria spp. | |
| cotoneaster | Cotoneaster apiculatus | |
| coyotebrush | Baccharis pilularis | |
| cycads | Cycas revoluta | |
| cypress, bald | Taxodium distichum | |
| cypress, Italian | Cupressus sempervirens | glauca |
| cypress, Japanese false | Chamaecyparis obtusa | oracis |
| cypress, Leyland | Cupressocyparis leylandii | |
| daylily | Narcissus spp. | King Alfred |
| daylily (sweet william) | Hemerocallis spp. | actes gold, bright yellow (hybrid), single gold (evergreen), wilson's yellow |
| delphinium | Delphinium spp. | magic fountain |
| dogwood | Cornus florida | |
| dogwood, American | Cornus sericea | flaviramea |
| douglas fir | Pseudotsuga menziesii | |
| duster miller | Senecio cineraria | |
| elm, drake | Ulmus parvifolia | |
| euonymus | Euonymus fortunei | argenteo-variegata, euzero-marginata, colorata, emerald gaily, emerald in gold, gold edge, gold princess, silver king, ticolet, vegetus |
| fan palm, European | Chamaerops humilis | |
| fan palm, Mexican | Washingtonia robusta | |
| fern (various) | Asparagus spp. | |
| fescue | Festuca glauca | |
| letterbush | Leucobae fontanesiana | rainbow |
| fig | Ficus retusa | robusta |
| fir, Fraser | Abies fraseri | |
| foxythia | Forsythia X intermedia | Arnold dwarf, bronzenis dwarf, lymwood gold, meadowlark |
| | Spring Glory | weeping |
| fuchsia | Fuchsia spp. | |
| gallium | Galium arduatum | |
| gardenia | Gardenia jasminoides | mystery, radicans |
| geranium | Pelargonium X hortorum | |
| gum | Eucalyptus citriodora | |
| hawthorn | Crataegus spp. | cockspur white, crimson cloud, enchantress, Jack Evans, Washington white |
| heather, twisted | Erica cinerea | Mediterranean pink |
| hemlock, Canada | Tsuga canadensis | |
| hibiscus | Hibiscus spp. | blue bird, brilliant, hula girl |
| holy | Ilex spp. | blue boy, blue girl |
| | Ilex X meserveae | burfordii, china girl, compacta, forsteri, helleri, Japanese northern beauty, needlepoint, Nellie R. Stevens |
| | Ilex X attenuata | Savannah |
| holy, Chinese | Ilex cornuta | |
| holy, Japanese | Ilex crenata | |
| holy, yaupon | Ilex vomitoria | |
| honeysuckle | Lonicera japonica | clayevii dwarf, halliana, tatarian Canadian white, zebeli red hosta |
| hosta | Hosta spp. | albo marginata |
| | Hosta japonica | |
| ice plant | Caryophytus edulis | |
| impatiens | Impatiens spp. | |
| iris | Iris spp. | dwarf blue, wedgewood |
| ivy, English | Hedera helix | bulgarica, thorndale |

| Common Name | Botanical Name | Tolerant Cultivars |
|-------------------------------|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| jasmine, Asian | Trachelospermum asiaticum | |
| juniper | Juniperus spp. | arcadia, armstrong, bar harbor |
| | Juniperus horizontalis | blue chip, blue pacific, blue rug |
| | Juniperus horizontalis | blue star, broadmoor, buffalo, Calgary carpet, emerald sea, emerald spreader, endora compacta, fruitland, green, gold lip, hetzi, hughes, Manhattan blue, parsoni, pilizeriana, plumosa, prince of Wales, procumbens dwarf, San Jose, sargent blue, sargent green, scandia, scopulorum moonloak, scopulorum styrocket, spartan |
| | Juniperus chinensis | lamariscifolia |
| | Juniperus sabina | weberi, youngslown, yukon belle |
| king palm | Archonophoenix cunninghamiana | |
| | Pittosporum tobira | |
| laurel, Australian | Kalmia latifolia | |
| laurel, mountain | Leucocoe fontanesiana | |
| leucotoe | Ligustrum japonicum | |
| ly, African | Agapanthus africanus | albus, Peter Pan |
| ly, African blue | Pieris japonica | Mt. Fire |
| ly of the valley | Liriope muscari | evergreen giant, lilac beauty, majestic, morrie white, variegata |
| lylart | Liriope spicata | |
| liriope, green | Acer japonicum | |
| maple, Japanese | Acer platanoides | |
| maple, Norway | Acer rubrum | |
| maple, red ¹ | Acer saccharinum | |
| maple, silver | Acer saccharum | |
| maple sugar ¹ | Taxus palustris | honeycomb, variegata, wheeleris dwarf |
| marigold | Philadelphus spp. | golden, snowflake double white |
| mock orange ¹ | Ophiopogon japonicus | |
| mondo grass | Portulaca grandiflora | sunnyside |
| moss rose | Lagerstroemia indica | laurel, langer, muskgoose, standard pink |
| myrtle, crepe | Mandarin domestica | compacta, nana |
| myrtle, wax | Narcissus spp. | |
| nandina | Quercus laurifolia | |
| narcissus | Quercus palustris | |
| oak, laurel | Quercus rubra | |
| oak, pin | Quercus virginiana | |
| oak, red | Quercus phellos | |
| oak, southern | Nerium oleander | hardy red, petite pink, Sister Agnes |
| oak, willow | Osteospermum fulcosum | willig |
| oleander | Pachysandra terminalis | |
| osteospermum | Coriaria selbiana | |
| pachysandra | Vinca viticelliana | |
| palm, bangalow | Bougainvillea glabra | Barbara karst |
| panicum grass | Prunus persica | |
| pantry | Vinca minor | |
| paper flower | Pelonia X hybrida | picot |
| peach ¹ | Phellina X Fraseri | |
| periwinkle, dwarf | Pieris japonica | |
| petunia | Pinus nigra | |
| phloxia, red tip | Pinus thunbergiana | |
| phloxia | Pinus taeda | |
| plum | Pinus koraiensis | |
| pine, Australian | Pinus mugo | |
| pine, Japanese black | Pinus strobus | |
| pine, loblolly | Pinus strobus | |
| pine, longleaf | Pinus palustris | |
| pine, mugo | Pinus mugo | |
| pine, Scotch | Pinus strobus | |
| pine, slash | Pinus strobus | |
| pine, Swiss mt. | Pinus strobus | |
| pine, Virginia | Pinus virginiana | |
| pine, white | Pinus strobus | |
| pineapple, guava ¹ | Ficus sellowiana | |
| plum | Pieris taiwanensis | |
| plum, Japanese ¹ | Pittosporum Japan | |
| potentilla | Potentilla nepalensis | abbotswood |
| | Potentilla fruticosa | |
| privet | Ligustrum japonicum | golden vicary, regal, lexanum, wax, yellow tipped |
| privet, glossy | Ligustrum lucidum | |
| pyracantha | Pyracantha koidzumii | gnome, islandi, victory |
| queen palm | Arecastum ramananzoffianum | |
| quince, Japanese ¹ | Rhododendron spp | album, Cunningham white, fashion, hardy, sym, purple gem, silver pink |
| rhododendron | | |

| Common Name | Botanical Name | Tolerant Cultivars |
|-------------------------|---------------------------|------------------------------------------------------------------------------------------------------|
| rhododendron carolina | Rhododendron carolinianum | |
| rhododendron calawba | Rhododendron catawbiense | |
| rockrose | Aralia spp. | snowflake |
| rhodie max (rosebay) | Rhododendron maximum | |
| rose ¹ | Rosa banksiae | lutea |
| rosemary ¹ | Rosmarinus officinalis | |
| rosemary, bog | Andromeda polifolia | nana |
| salvia | Salvia farinacea | rhea |
| sedum | Sedum spurium | dragon blood red, red carpet, yellow |
| snagdragon | Anemithium spp. | |
| sourwood | Oxydendrum arboreum | |
| spiraea | Astibe X arendsii | fanall |
| spiraea | Spiraea spp. | Anthony waterer red, daichica, froebel pink, goldenflame red, snowmound white, Van Houtte white |
| spiraea, garland | Spiraea spp. | |
| spruce, black hills | Picea glauca | |
| spruce, Colorado blue | Picea pungens | |
| spruce, norway | Picea abies | |
| spruce, white | Picea glauca | conica |
| sweetflag, grassy-le | Liquidambar styraciflua | |
| sweetgum | Pistacia occidentalis | |
| sycamore | Asparagus virginicus | |
| tree fern (liri lem) | Gelsomium sempervirens | |
| trumpet flower, evening | Tulip spp. | apeldoorn |
| tulip | Lantana sellowiana | |
| verbena, shrub | Verbena spp. | St. Paul |
| vervain | Viburnum spp. | American cranberry bush, arrowwood, common snowball, European cranberry bush, Inden, Mohican, Wright |
| viburnum | Vinca minor | |
| vinca (periwinkle) | Trachycarpus fortunei | |
| windmill palm | Xylopa congestum | |
| xylosma | Achillea spp. | |
| yarrow | Ilex vomitoria | dwarf |
| yaupon | Taxus cuspidata | densiformis |
| yew | Taxus X media | |

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC
EPA accepted 03/24/03
RD 091912