



**Spread It and Forget It Fertilizer with Dimension® 0.15%**

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

This bag covers 11,350 sq ft

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, this product may be applied only by commercial applicators at no more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Active Ingredient dithiopyr, 3,5-pyridinedicarboxylic acid-(2-difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester 0.15%  
Other Ingredients 99.85%

Total 100.0%

This product contains 0.075 pound of the active ingredient dithiopyr per 50 pound bag

**Keep Out of Reach of Children**

**CAUTION**

- |  |  |   |
|--|--|---|
| EPA EST. NO. 87245-CAN-1 (Lot begins 2*) | EPA Reg. No. 62719-504-84886               | EPA EST. NO. 49991-TX-2 (Lot begins TX-2) |
| "If" lot begins "2", PRODUCT OF CANADA   | EPA EST. NO. 75527-PA-12 (Lot ends PA-012) | EPA EST. NO. 89135-AZ-1 (Lot begins 4)    |
| EPA EST. NO. 41124-WH-1 (Lot begins W)   | EPA EST. NO. 69759-OR-2 (Lot begins MAS)   | EPA EST. NO. 35512-FL-2 (Lot ends GF)     |
| EPA EST. NO. 57131-WH-1 (Lot begins EC)  | EPA EST. NO. 41124-OH-1 (Lot begins O)     | EPA EST. NO. 62757-FL-1 (Lot begins S)    |
| EPA EST. NO. 82757-OH-1 (Lot begins MF)  | EPA EST. NO. 82757-MA-2 (Lot begins H)     |   |

**Net Weight 50 LBS. (22.68 Kg)**

**Precautionary Statements**

**Hazards to Humans and Domestic Animals**

**Causes Eye Irritation • Harmful If Inhaled**

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

**First Aid**

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

**Environmental Hazards**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**Directions for Use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Storage and Disposal**

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.  
**Pesticide Storage:** Store this product only in its original container in a dry, cool, secured storage area.  
**Pesticide Disposal:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.  
**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Use Directions for Turf**

This product provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes.

When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

**Use Precautions for Turf**

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may increase the chances of turf injury. Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed.

Cool-Season Grasses	Warm-Season Grasses	
bentgrass, creeping <sup>†</sup>	Agrostis palustris	bahiagrass
bluegrass, Kentucky	Poa pratensis	bermudagrass
fescue, fine <sup>††</sup>	Festuca rubra	buffalograss <sup>†††</sup>
fescue, tall	Festuca arundinacea	carpetgrass
ryegrass, perennial	Lolium perenne	centipedegrass
		kikuyugrass
		St. Augustinegrass
		zoysiagrass
		Zoysia japonica
		Paspalum rotatum
		Cynodon dactylon
		Buchloe dactyloides
		Axonopus affinis
		Eremochloa ophiuroides
		Pennisetum clandestinum
		Stenotaphrum secundatum

<sup>†</sup> Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colonial bentgrass (Agrostis tenuis).

<sup>††</sup> Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlantia', 'Banner', 'Beauty', 'Bigarf', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivala', 'Jamestown', 'Kokoi', 'Majenta', 'Mary', 'Penlawn', 'Tamara', 'Tajana', 'Waldorf', and 'Wilding'. Not all varieties of fine fescue have been tested.

<sup>†††</sup> Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

**Note:** It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

**Precautions for Reseeding, Overseeding, or Sprigging**

- Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 7.7 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses.
- When reseeding or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation and fertilization.
- For best overseeding results, use mechanical or power seeding equipment (silt seeders) designed to provide good contact between soil and seed.

**Precautions for Control of Crabgrass**

Early postemergence weed control with this product is limited to crabgrass. Postemergence application effectively controls crabgrass, but only if application is made prior to the 3-leaf stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

**Other Use Precautions**

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.
- Do not graze livestock or feed forage cut from areas treated with this product.

**Application Directions for Turf**

**Tips for Improved Weed Control**

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Use of split (half-rate) applications spaced 5 - 10 weeks apart may provide improved weed control.

**Maximum Use Rates (Applies to all Turfgrass Uses)**

- Do not apply more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per application or more than 1000 lb (1.5 lb active ingredient) per acre (23.1 lb per 1000 sq ft) per year if using split or sequential applications.
- In the state of New York, this product may be applied only by commercial applicators at no more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

**37-0-5**

**GUARANTEED ANALYSIS**

Total Nitrogen (N) ..... 37.00%  
37.00% Urea Nitrogen\*  
Soluble Potash (K<sub>2</sub>O) ..... 5.00%

DERIVED FROM: Polymer Coated Urea, Urea and Muriate of Potash \*31.6% Slowly Available Urea Nitrogen from Polymer Coated Urea

WARNING: THIS PRODUCT CONTAINS CHEMICALS KNOWN TO THE STATE OF CALIFORNIA TO CAUSE CANCER OR BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM. Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP" for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/bios/nonpoint/gf/bmp07.pdf>.

Florida Applications: For application to urban turf (non-agricultural land planted in closely mowed, managed grasses except golf courses, parks and athletic fields), follow "Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries" available at [http://www.dep.state.fl.us/central/Home/Meetings/Training/FLGreenBMP\\_Book\\_final.pdf](http://www.dep.state.fl.us/central/Home/Meetings/Training/FLGreenBMP_Book_final.pdf)

Product No.6000272

F1828

**GENERAL APPLICATION GUIDELINES**

Rate	lbs. Product per 1,000 sq ft	lbs. Nitrogen per 1,000 sq ft	lbs. A.I. per acre	
Low	5.4	2.0	0.35	
Medium	6.4	2.4	0.42	
High	7.6	2.8	0.50	
Spreader Setting	AccuPro2000	Bigfoot	Lesco Rotary	Spyker
Low	P	O	28	7
Medium	R	P	30	7.5
High	T	Q	19 1/2*	8

\*These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product. \*Use lower setting and spread in 2 directions to achieve high rate with this spreader.



This product contains Duraton CR, a registered trademark of Agrium, Inc.



**GUARANTEED BY:**  
Agrium Advanced Technologies (U.S.) Inc.  
2915 Rocky Mountain Avenue  
Loveland, CO 80538

## Control of Crabgrass

### Preemergence and Early Postemergence Control

- This product provides "preemergence" control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass.
- This product also provides "early postemergence" control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergence crabgrass control is a 2 - 4 week longer period of time (window of application) during which crabgrass control can be achieved.

### Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described in the following use programs.

## Spring Applications

For applications made in the spring or early summer, use an appropriate application rate corresponding to one of the three control programs listed in Table 1. Specific rate recommendations are based on the user's location (Region), the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

**Use Program 1** for preemergence control at sites where turfgrass is cut relatively high (e.g., homeowner lawns). This program provides 3 - 5 months of preemergence crabgrass control. This program will also provide early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., homeowner lawns).

**Use Program 2** for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., residential lawns).

**Use Program 3** for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively low (e.g., golf fairways).

Table 1 Recommended Use Rates

Region	Application Rates	Program 1	Program 2	Program 3
<b>All states, except NY, and parts of states not included in transition, south, coastal south or west.</b>	lb/acre	127	167	253
	lb/1000 sq ft	2.9	3.8	5.8
	lb ai/acre	0.19	0.25	0.38
<b>Transition:</b> DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, & RI	lb/acre	167	253	333
	lb/1000 sq ft	3.8	5.8	7.7
	lb ai/acre	0.25	0.38	0.5
<b>South:</b> AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX	lb/acre	253	167 + 167	207 + 207
	lb/1000 sq ft	5.8	3.8 + 3.8 <sup>†</sup>	4.7 + 4.7 <sup>†</sup>
	lb ai/acre	0.38	0.25 + 0.25	0.31 + 0.31
<b>Coastal South:</b> HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX	lb/acre	167 + 167	207 + 207	253 + 253
	lb/1000 sq ft	3.8 + 3.8 <sup>†</sup>	4.7 + 4.7 <sup>†</sup>	5.8 + 5.8 <sup>†</sup>
	lb ai/acre	0.25 + 0.25	0.31 + 0.31	0.38 + 0.38
<b>West:</b> AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	lb/acre	127 + 167	167 + 253	167 + 167
	lb/1000 sq ft	2.9 + 3.8	3.8 + 5.8	3.8 + 3.8 <sup>†</sup>
	lb ai/acre	0.19 + 0.25	0.25 + 0.38	0.25 + 0.25

<sup>†</sup> Split applications may be made 5 - 10 weeks apart.

**Use of Split or Sequential Applications:** Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with the rates in Table 1 being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application, provided the maximum application rate per year is not exceeded. Refer to the "Maximum Use Rates" section above and separate rate restrictions for the state of New York.

## Fall Applications

This product can also be applied in the late summer through early fall (late August through November) at use rates listed in Table 1 for "Program 3" to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for seasonal control of crabgrass.

## Weeds Controlled or Suppressed

### Spring Applications

When applied preemergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds at application rates given in the Table 1 above:

Grasses			
barley	<i>Hordeum</i> spp.	goosegrass	<i>Elysiue indica</i>
barnyardgrass	<i>Echinochloa crus-galli</i>	foxtail, green	<i>Setaria verti</i>
bluegrass, annual	<i>Poa annua</i>	foxtail, yellow	<i>Setaria faberi</i>
brome	<i>Bromus</i> spp.	kikuyugrass †	<i>Pennisetum clandestinum</i>
crabgrass, large	<i>Digitaria sanguinalis</i>	oats, wild	<i>Avena fatua</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>	ryegrass (annual & perennial)	<i>Lolium</i> spp.
crabgrass, Southern	<i>Digitaria ciliaris</i>	sandbur	<i>Cenchrus</i> spp.
crowfootgrass †	<i>Dactyloctenium aegyptium</i>	smutgrass	<i>Sporobolus indicus</i>
dallisgrass (seeding)	<i>Paspalum dilatatum</i>		
Broadleaf Weeds			
bittercross †	<i>Cardamine</i> spp.	pigweed, redroot	<i>Amaranthus retroflexus</i>
carpetweed	<i>Mollugo verticillata</i>	parsley-piert †	<i>Alchemilla arvensis</i>
chickweed †	<i>Stellaria</i> spp.	purslane, common	<i>Portulaca oleracea</i>
geranium, Carolina †	<i>Geranium carolinianum</i>	rocket, London	<i>Sisymbrium ino</i>
herbit	<i>Lamium</i> spp.	shepherdspurse	<i>Capsella bursa-pastoris</i>
knockweed, prostrate	<i>Polygonum aviculare</i>	speedwell, com †	<i>Veronica arvensis</i>
lespedeza, common †	<i>Lespedeza striata</i>	spurge, garden	<i>Euphorbia hirta</i>
marehail	<i>Conyza canadensis</i>	spurge, prostrate	<i>Euphorbia humistrata</i>
medic, black	<i>Medicago lupulina</i>	spurge, spotted	<i>Euphorbia maculata</i>
mustard	<i>Brassica</i> spp.	woodsorrel, creeping	<i>Oxalis corniculata</i>
oxalis, buttercup	<i>Oxalis pes-caprae</i>	woodsorrel, yellow	<i>Oxalis stricta</i>
pineappleweed †	<i>Matricaria matricarioides</i>		

<sup>†</sup> Suppression only

### Fall Applications

When applied preemergence (prior to germination) in late summer through early fall (late August through November) this product will control or suppress the following weeds in addition to providing preemergence control of crabgrass in the following spring:

bluegrass, annual	<i>Poa annua</i>	geranium, Carolina	<i>Geranium carolinianum</i>
bittercross	<i>Cardamine</i> spp.	parsley-piert	<i>Alchemilla arvensis</i>
chickweed	<i>Stellaria</i> spp.	pineappleweed	<i>Matricaria matricarioides</i>
herbit	<i>Lamium</i> spp.	shepherdspurse	<i>Capsella bursa-pastoris</i>

### Application Instructions and Equipment

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

## Use Directions for Landscape Ornamentals

This product provides preemergence control or suppression of listed annual grass and broadleaf weeds in plantings of ornamental plants listed on this label. This product may be applied in ornamental gardens, parks, golf courses and residential areas where ornamental plants are grown for aesthetic purposes. Apply this product with a properly calibrated spreader that will provide uniform particle distribution. Follow instructions in the "Application Instructions and Equipment" section above.

### Use Rates

This product may be applied as a single application or split application. Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. To make a split application, divide the rate in the following table into separate applications made 5-10 weeks apart. Sequential applications may be made in the late summer following a spring application or in the spring after a fall application, provided maximum application rates per year given below are not exceeded.

Pounds of Product/Acre	Pounds of Product/1000 sq ft	Pounds of Product/100 sq ft <sup>†</sup>
333	7.7	0.77 (12.3 oz)

<sup>†</sup> Rate based on an area of 100 sq ft for treatment of small areas.

### Maximum Use Rates

- Do not apply more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per application or more than 1000 lb (1.5 lb active ingredient) per acre (23.1 lb per 1000 sq ft) per year if using split or sequential applications.
- In the state of New York, this product may be applied only by commercial applicators at no more than 333 lb (0.5 lb active ingredient) per acre (7.7 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

### Application Timing and Recommendations

This product is effective as a preemergence herbicide, but will not control established weeds. Applications to mulched areas or bare ground must be made prior to weed seed germination. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Existing weedy vegetation may be controlled by hand weeding, cultivation or using postemergence herbicides prior to application.

### Tips for Improved Weed Control

- The herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Avoid disturbance of treated areas. Loss of weed control may result if the treated soil surface is disturbed by soil mixing or tillage.

### Use Precautions

- Apply this product only to established ornamentals.
- Do not apply this product directly to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress resulting from drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides, or damage due to insects, nematodes or disease.
- Do not allow foilar damage to ornamentals, irrigate immediately after application to remove any herbicide granules adhering to foliage.
- Do not apply under conditions that would affect uniformity of application or distribution on the soil surface. Uneven product distribution will result in uneven weed control. Application under windy conditions can result in uneven distribution or cause herbicide granules to drift from the intended treatment area.
- Do not graze livestock or feed forage cut from areas treated with this product.

### Weeds Controlled or Suppressed

Used as directed, this product will control crabgrass and control or suppress other listed grass and broadleaf weeds when applied prior to their germination. Refer to "Use Directions for Turf" section for a complete listing of weeds controlled or suppressed. This product will not control established broadleaf weeds or grasses, except for crabgrass in early stages of development. The area to be treated should be free of weeds prior to application.

### Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to directions in this label. However, this product has not been tested on all ornamental species, all cultivars of species on this label, or under all possible growing conditions. For species or cultivars not listed, it is recommended that a few plants in a limited area be treated and observed for tolerance under local growing conditions prior to large-scale use.

Common Name	Botanical Name	Tolerant Cultivars
abelia, dwarf	<i>Abelia X grandiflora</i>	nana
ajuga	<i>Ajuga reptans</i>	bronze
	<i>Ajuga renevenensis</i>	bronze beauty
almond, flowering	<i>Prunus glandulosa</i>	
apple <sup>†</sup>	<i>Malus pumila</i>	
arbutovite	<i>Thuja occidentalis</i>	nigra, pyramidalis, amargah, tochny, woodwards
arbutovite, golden	<i>Thuja orientalis</i>	
aster, Chinese	<i>Callistephus chinensis</i>	dwarf queen
ash, green	<i>Fraxinus pennsylvanica</i>	
ash, mountain	<i>Sorbus aucuparia</i>	
ash, purple	<i>Fraxinus americana</i>	
azalea	<i>Rhododendron</i> spp.	brilliant, buccaneer, caror, chimes (belgian), elise lee, exbury, lashion, hardizer beauty, hershey red, ligase, hincorinson, holland (hybrid), marion lee, northern lights, orange cup, orchid lights, snow, southern charm
azalea, flame	<i>Rhododendron calendulaceum</i>	
azalea, kishima		
bamboo, heavenly		
barberry	<i>Berberis thunbergii</i>	aurca, dwarf pigmy, green, kobold, pygmy red, rose glow
barberry, purple		atrapurpurea
basket flower	<i>Gaillardia grandiflora</i>	
bearberry (common)	<i>Arctostaphylos uva-ursi</i>	Massachusetts
bee balm	<i>Monarda didyma</i>	
begonia	<i>Begonia</i> spp.	
berch, river	<i>Betula nigra</i>	
blackeyed Susan	<i>Rudbeckia hirta</i>	goldstrum

Common Name	Botanical Name	Tolerant Cultivars
blanket flower	Gaillardia spp.	
blueberry 1	Vaccinium spp.	bluecrop, blue ray, jersey, north blue, northland
botchbrush	Callistemon ciliatus	
boxwood, Japanese		japonica
boxwood, weller	Buxus sempervirens	
broom	Cytisus spp.	moonlight
	Genista plicata	Vancouver gold
bugle carpet		
camellia	Camellia japonica	debutante, matholana supreme, chansonette
	Camellia sasanqua	
candy tuft	Iberis spp.	snow white
carex, variegated	Carex	
cedar, red	Juniperus virginiana	
celosia	Celosia spp.	
centaurea	Centaurea montana	
cockscomb, plumosa	Celosia cristata	scarlet plumosa
colerus	Coleus blumei	red kewpie
columbine	Aquilegia spp.	
copper leaf	Acalypha wilkesiana	
coreopsis	Coreopsis spp.	moonbeam
corn flower	Centaurea spp.	
cornnester	Cotoneaster apiculatus	
corylebrush	Baccharis pilularis	
cycads	Cycas revoluta	
cypress, bald	Taxodium distichum	
cypress, Italian	Cupressus sempervirens	glauca
cypress, Japanese false	Chamaecyparis obtusa	gracilis
cypress, Leyland	Cupressocyparis leylandii	
daffodil	Narcissus spp.	King Alfred
daylily	Hemerocallis spp.	aztec gold, bright yellow (hybrid), single gold (evergreen), wiltons yellow
dianthus (sweet william)	Dianthus spp.	
delphinium	Delphinium spp.	magic fountain
dogwood	Cornus florida	
dogwood, American	Cornus sericea	flaviramea
douglas fir	Pseudotsuga menziesii	
dusty miller	Senecio cineraria	
elm, drake	Ulmus parvifolia	
eunonymus	Eunonymus fortunei	argenteo-variegata, aureo-marginata, colorata, emerald gaiety, emerald én gold, gold edge, gold princess, silver king, tricolor, vegetus
fan palm, European	Chamaerops humilis	
fan palm, Mexican	Washingtonia robusta	
firm (various)	Asparagus spp.	
fescue	Festuca glauca	
fetterbush	Leucothoe fontanesiana	rainbow
ficus	Ficus retusa	nitida
fir Fraser	Abies fraseri	
forsythia	Forsythia X intermedia	Arnold dwarf, bronensis dwarf, lymwood gold, meadowlark
	Spring Glory	weeping
fuchsia	Fuchsia spp.	
gallium	Galium ordoatum	
gardenia	Gardenia jasminoides	mystery, radicans
geranium	Pelargonium X hortorum	
gum	Eucalyptus citrodora	
hawthorn	Crataegus spp.	cockspur white, crimson cloud, enchantment, Jack Evans, Washington white
heather, twisted	Erica cinerea	Mediterranean pink
hemlock, Canada	Tsuga canadensis	
hibiscus	Hibiscus spp.	blue bird, brilliant, hula girl
holly	Ilex spp.	blue boy, blue girl
	Ilex X meserveae	burford, china girl, compacta, forsteri, helleine, Japanese northern beauty, needpoint, Nellie R. Stevens
	Ilex X attenuata	Savannah
holly, Chinese	Ilex cornuta	
holly, Japanese	Ilex crenata	
holly, yaupon	Ilex vomifolia	
honeysuckle	Lonicera japonica	clayvii dwarf, halliana, tatarian Canadian white, zebell red hosta
hosta	Hosta spp.	albo marginata
	Hosta lancifolia	
ice plant	Carportrotus edulis	
impatiens	Impatiens spp.	
iris	Iris spp.	dwarf blue, webdewood
ivy, English	Hedera helix	bulgaria, thornlate

Common Name	Botanical Name	Tolerant Cultivars
jasmine, Asian	Trachelospermum asiaticum	
juniper	Juniperus spp.	arcadia, armstrong, bar harbor
	Juniperus horizontalis	blue chip, blue pacific, blue rug
	Juniperus horizontalis	blue star, broadmoor, buffalo, Calgary carpet, emerald sea, emerald spreader, endora compacta, itulifand, green, gold tip, hetzi, hughes, Manhattan blue, parsoni, pitzeriana, plumosa, prince of Wales, procumbens dwarf, San Jose, sargent blue, sargent green, scandia, scopulorum moon glow, scopulorum skyrocket, spartan
	Juniperus chinensis	tamariscobila
	Juniperus sabinna	weberi, youngstown, yukon belle
king palm	Archontophoenix cunninghamiana	
laurel, Australian	Pittosporum tobira	
laurel, mountain	Kalmia latifolia	
leucothoe	Leucothoe fontanesiana	
ligustrum, Japanese	Ligustrum japonicum	
lily, African	Agapanthus africanus	albus, Peter Pan
lily, African blue	Renele japonica	Mt. Fm
lily of the valley	Lilium muscari	evergreen giant, lilac beauty, majestic, moroze white, variegata
lylirt	Liriodendron tulipifera	
liriodendron	Liriodendron tulipifera	
liriodendron, green	Acer japonicum	
maple, Japanese	Acer japonicum	
maple, Norway	Acer dasycarpum	
maple, red 1	Acer rubrum	
maple, silver	Acer saccharinum	
maple, sugar 1	Acer saccharum	
marigold	Tagetes patula	honeycomb, variegata, wheel/iris dwarf
mock orange 1	Philadelphus spp.	golden, snowflake double white
mondo grass	Ophiopogon japonicus	
moss rose	Portulaca grandiflora	sunrise
myrtle, crape	Lagerströmia indica	laurel, lanser, muskoguee, standard pink
myrtle, wax	Myrica cerifera	
nandina	Nandina domestica	compacta, nana
narcissus	Narcissus spp.	
oak, laurel	Quercus laurifolia	
oak, pn	Quercus palustris	
oak, red	Quercus rubra	
oak, southern	Quercus virginiana	
oak, willow	Quercus phellos	
oleander	Nerium oleander	hardy red, petite pink, Sister Agnes
ostelepernum	Ostelepernum fruticosum	wiflag
pachysandra	Pachysandra terminalis	
palm, bangalow	Coradorea zebrana	
pampas grass	Viola X wittrockiana	
pansy	Bougainvillea glabra	Barbara karst
paper flower	Prunus persica	
peach 1		
petunick, dwarf	Vinca minor	
petunia	Petunia X hybrida	picot
phloxia, red tip	Phloxia X fraseri	
piers	Pieris japonica	
pine, Australian	Pinus nigra	
pine, Japanese black	Pinus thunbergiana	
pine, loblolly	Pinus taeda	
pine, longleaf	Pinus palustris	
pine, mugo	Pinus mugo	
pine, Scotch	Pinus sylvestris	
pine, slash	Pinus elliotii	
pine, Swiss mt.	Pinus mugo	
pine, Virginia	Pinus virginiana	
pine, white	Pinus strobus	
pineapple, guava 1	Frejya zellowiana	
piers	Pieris taiwanensis	
pittosporum, Japan	Pittosporum japonicum	
potentilla	Potentilla nepalensis	abbotswood
	Potentilla fruticosa	
privet	Ligustrum japonicum	golden vicary, regal, lexanum, wax, yellow tipped
privet, glossy	Ligustrum lucidum	
pyracantha	Pyracantha coccinifera	gnome, island, victory
queen palm	Arecastrum zamantioffianum	
quince, Japanese 1		
rhododendron	Rhododendron spp.	albus, Cunningham white, fashion, hardy, pm, purple gem, silvery pink

Common Name	Botanical Name	Tolerant Cultivars
rhododendron, carolina	Rhododendron carolinianum	
rhododendron, catarata	Rhododendron catawbiense	
rockrose	Azalea spp.	snowcap
rhodie max (rosebay)	Rhododendron maximum	
rose 1	Rosa banksiae	lutea
rosemary 1	Rosmarinus officinalis	
rosemary, bog	Andromeda polifolia	nana
salvia	Salvia farinacea	rhea
sedum	Sedum spurium	dragon blood red, red carpet, yellow
snadragon	Anemone spp.	
sourwood	Corydalis arborea	
spiraea	Asilbe X arendsi	fanall
spiraea	Spiraea spp.	Anthony waterer red, doichica, freebel pink, goldenfame red, snowmound white, Van Houtte white
spiraea, garland	Spiraea spp.	
spruce, black hills	Picea glauca	
spruce, Colorado blue	Picea pungens	
spruce, norway	Picea abies	
spruce, white	Picea glauca	corica
sweetflag, grassy-le	Liquidambar styraciflua	
sweetgum	Liquidambar styraciflua	
sycamore	Platanus occidentalis	
tee fern (iki fern)	Asparagus virgatus	
tumpet flower, evening	Gelsemium sempervirens	
tulip	Tulip spp.	apeldoorn
verbena, shrub	Lantana zellowiana	
vervain	Verbena spp.	St. Paul
virburnum	Virburnum spp.	American cranberry bush, arrowwood, common snowball, European cranberry bush, linden, Mahican, Wright
vinca (periwinkle)	Vinca minor	
windmill palm	Trachycarpus fortunei	
xyloma	Xyloma congestum	
yarow	Achillea spp.	
yaupon	Ilex vomifolia	dwarf
yew	Taxus cuspidata	densiformis
	Taxus X media	

1 Ornamental species only. Do not use on food producing trees and ornamentals.

Terms and Conditions of Use  
 If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer  
 Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use  
 It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, borndees, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks shall be assumed by Buyer.

Limitation of Remedies  
 To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:  
 1. Refund of purchase price paid by buyer or user for product bought, or  
 2. Replacement of amount of product used.

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.  
 The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

\*Trademark of Dow AgroSciences LLC  
 EPA accepted 0324/03  
 RD 080613