

All-Season

CRABGRASS CONTROL PLUS

With Prodiamine

Fertilizer that builds thick, green lawns. Stops many weed grasses before they grow.




ACTIVE INGREDIENTS:		
Prodiamine: (CAS 26091-21-2)	0.29%
INERT INGREDIENTS:	99.71%
	Total	100.00%

EPA Reg. No. 32802-70
EPA Est. No. 32802-MO-1 (B)

16·0·4

Covers 5,000 sq. ft. Net Weight: 15 lbs. (6.8 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR FURTHER
PRECAUTIONS AND FIRST AID INFORMATION.



Guaranteed Analysis

Total Nitrogen (N)	16%
16.00% Urea Nitrogen	
Soluble Phosphorus (P ₂ O ₅)	4%
Derived From: Monoammonium Phosphate Urea	
Calcium (Ca), not more than 6.00%	0.31%

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.espfco.org/metals.htm>

Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS: This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

FIRST AID	
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Hot line number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

General Information

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens). It controls susceptible weeds by inhibiting weed seeds germination and root development. When used according to directions, it will control: bermudagrass; bluegrass, annual (Poa annual); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed);

lamb'squarters; knotweed; Kochia; lambsquarters, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass²; shepherd's purse²; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹ If heavy annuals a single application of 5.2 to 11.9 pounds/1000 square feet will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 5.2 to 7.9 pounds/1000 square feet, followed after 60 to 90 days by a second application at rates that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

² For this weed, apply in late summer, fall or winter prior to germination.
³ Suppression only.

When And How To Apply

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days after application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



Approximate Date

- After May 30
- After May 10
- After April 20
- After March 20
- January 1 to March 20

Turf Species	Minimum Rate	Maximum Annual Rate ¹
Bahiagrass, Bermudagrass ²	2.75 lbs / 1000 sq ft	11.9 lbs / 1000 sq ft per year
Centipedegrass, Zoysiagrass		
Seashore Paspalum		
St. Augustinegrass		
Tall Fescue (including turf-type)	2.75 lbs / 1000 sq ft	7.9 lbs / 1000 sq ft per year
Buffalograss		
Kentucky Bluegrass		
Perennial Ryegrass		
Creeping Red Fescue	2.75 lbs / 1000 sq ft	6 lbs / 1000 sq ft per year
Creeping Bentgrass	2.75 lbs / 1000 sq ft	5.2 lbs / 1000 sq ft per year

¹ These are the maximum rates per calendar year.

² May be used on newly sodged or plugged Bermudagrass at rates not to exceed 4.1 pounds/1000 square feet. Newly sodged or plugged Bermudagrass that is not sodged may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sodging situations.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
2.75 pounds / 1000 sq ft	4	4	4
5.2 pounds / 1000 sq ft	5	4	4
6 pounds / 1000 sq ft	6	5	5
6.3 pounds / 1000 sq ft	—	6	6
7.9 pounds / 1000 sq ft	—	7	7
9 pounds / 1000 sq ft	—	—	9
10.3 pounds / 1000 sq ft	—	—	10
11.9 pounds / 1000 sq ft	—	—	12

Spreader Settings

Broadcast Spreader	Drop Spreader
Agri-Fab	Agri-Fab
Ames	Ames
Ames Deluxe 2in1	Ames Deluxe 2in1
Craftsman	Craftsman
Earthway	Earthway
Plant Mate	Plant Mate
Precision GreenLine	Precision GreenLine
Precision Red Devil	Precision Red Devil
Precision Pro	Precision Pro
Scotts Speedy Green	Scotts Accu Green
Scotts Easy Green	Scotts Even Green
Scotts Basic	Taylor
Scotts Lawn Pro	
Scotts Deluxe EdgeGuard	
Taylor	

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

IF EMPTY: Do not reuse this container. Place in trash or offer for recycling if available.

IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

WARRANTY AND DISCLAIMER: Manufacturer/seller warrants that this product conforms to the ingredient statement on the label, and that the product is reasonably suited for the labeled use when applied according to directions. No other warranty, expressed or implied, is made concerning the merchantability or fitness for a particular purpose. Manufacturer/seller's liability shall be limited to the purchase price of the product. In no case shall manufacturer/seller be liable for consequential or incidental damages or loss. Buyer and user assumes all risks and liability of use, storage and/or handling of this product not in accordance with the terms of this label.



PKR-UY080
FGR07010 C70-00872 10/08

Manufactured and Guaranteed By



Howard Johnson's Enterprises, Inc.
9575 S 60th Street, Franklin, WI 53132

© Copyright 2008
Howard Johnson's Enterprises, Inc.