


vegalab

NON-TOXIC
Environmentally Safe



MILDEW CONTROL

ALL NATURAL MILDEW CONTROL

LIQUID CONCENTRATE

ACTIVE INGREDIENT:	
Geraniol (CAS# 106-24-1)	16.7%
INERT INGREDIENTS:*	83.3%
TOTAL:	100.0%

*Water (CAS# 7732-18-5), Isopropyl alcohol (CAS# 67-63-0)

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See Back Panel for Precautionary Statements


Manufactured by: Vegalab LLC
1201 North Orange Street, Suite 7233
Wilmington, DE 19801-1186

NET CONTENTS: 32 fl. ozs. (946 ml)

Naturally Controls Downey and Powdery Mildew

Mildew Control is a fungicidal agent derived from organic plant extracts. It is applied as a foliar spray and is effective against downey and powdery mildew.

Mildew Control uses natural extracts of Geraniol as a fungicide. Mildew growth cycles begin with spore germination at the plant surface. After docking at the surface, germ tubes begin to grow and feed from plant cell nutrients. The two-fold prevention and healing effect works by inhibiting spore germination, germ tube growth, and by destroying the pathogen cell. Geraniol induces an unbalance of K⁺ ions inside the mycelium and inhibits spore germination and mycelium proliferation. It enters the pathogen cell, causing cell lyses.



vegalab



MILDEW CONTROL

ALL NATURAL MILDEW CONTROL

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. **IF INHALED** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. **IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information call the National Pesticides Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a week. During other times, call the poison control center 1-800-222-1222.

- Effective against *Podospaera fusca*, the powdery mildew found on cucumbers, luffas, melons (including watermelons), and squashes (including pumpkins).
- Effective against *Erysiphe necator* (*Uncinula necator*), the powdery mildew endemic to grapes.
- Effective against *Podospaera leucotricha*, the powdery mildew that affects both apples and pears.
- Effective against *Podospaera aphanis*, the powdery mildew commonly found on rosaceae, such as blackberries, raspberries, and roses.
- Effective against *Oidium lycopersicum*, the powdery mildew of tomatoes.
- Effective against powdery mildew on ornamental and lawn and turf.

DIRECTIONS FOR USE

Mix Rate: The mix rate for Mildew Control concentrate is 0.13 fl. oz. (3/4 tsp.) or 3.8 mL per one gallon (3.79 L) of water. SHAKE OR MIX THOROUGHLY!

Application Table

ALL CROPS, EXCEPT STRAWBERRY*

Mix Rates	Dilution Rate	Foliar Spray	Timing & Frequency
0.13 fl. oz. (3.8 mL)/ 1 gal.	1:1,000	0.15 fl. oz. (4.5 mL)/1,000 ft ²	Low Severity disease outbreak
		7.03 fl. oz. (208 mL)/Acre 16.9 fl. oz. (500 mL)/Ha	
12.8 fl. oz. (379 mL)/ 100 gal.		0.24 fl. oz. (7 mL)/1,000 ft ²	High Severity disease outbreak
		10.56 fl. oz. (312.5 mL)/Acre 25.36 fl. oz. (750 mL)/Ha	

*Not recommended for use on strawberries, as it may cause phytotoxicity to flowers.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This product has not been registered by the United States Environmental Protection Agency. Vegalab LLC represents that this product qualifies for exemption from registration under FIFRA 25(b). ESL25b04 REV091113

