

vegalab



SPIDER MITE CONTROL

ALL NATURAL ACARICIDE

FIRST AID

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. IF INHALED • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice. IF IN EYES • Hold eye open and rinse slovily and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue mining eye. • Call a poison control center or doctor for treatment advice. IF ON SKIN OR CLOTHING - Take off contaminated clothing. • Rinse skin immediately with plently of water for 15-20 minutes. • Call a poison control center or doctor for treatment.

HOT LINE NUMBER: Have the product

control center of doctor for treatment HOT LINE NUMBER: Have the product container or label with you when calling a poison control center of doctor, or going for treatment. For emergency information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific time (PT), seven days a vieek. During other times, call the poison control center 1-800-222-1222.

EFFICACY

- Effective against all spider mites and on any crop, lawn, ornamental, or turf.
- · Tested on Tetranychus Urticae, which attacks beans, corn, peppers, potatoes, strawberries, tomatoes, and ornamentals such as roses.
- Effective as insect repellent, broadly recommended, and used in various fields, such as forestry, horticulture, lawn care, and organic farming.

DIRECTIONS FOR USE

Mix Rate: The mix rate for Spider Mite Control is 1.28 fl. oz. (37.9 mL) per one gallon of water that may only be used as a foliar spray. SHAKE OR MIX THOROUGHLY!

Application Table

Plant Type	Mix Rates	Dilution Rate	Foliar Spray	Time To Use
All crops, lawns, or ornamentals	1.28 fl. oz. (37.9 mL)/1 gal. 1 gal. (3.79 L)/100 gal.	1:100	15.3 fl. oz. (452 mL)/1,000 ft ² 5.2 gal. (19.7 L)/Acre 12.5 gal. (47.3 L)/Ha	During time of spider mite occurrence

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows. Empty the remaining contents into application equipment or a container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Darin for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. This product has not been registered by the United States Environmental Protection Agency, Vegalab LLC represents that this product qualifies for exemption from registration under FIFRA 25(b).

ESL25b07 REV090713